

Review (Mid Term #2)

Ex. Let $f(x) = \frac{1}{1-4x}$. Find the equation of tangent at $x=0$.

$$\bullet \quad f(x) = \frac{1}{1-4x} \Rightarrow f(0) = \frac{1}{1-4(0)} = 1.$$

So, $(0, 1)$ point is on the graph of $f(x)$.

We are seeking the tangent of $f(x)$ at $(0, 1)$ point.

• Slope of $f(x)$ at $x=0$ is:

$$f'(x) = -\frac{1}{(1-4x)^2} \cdot (-4) = \frac{4}{(1-4x)^2}$$

$$f'(0) = \frac{4}{(1-4(0))^2} = \frac{4}{1} = 4.$$

So the equation of the tangent passing through $(0, 1)$ with slope at $x=0$ is 4 is given by:

$$y - 1 = 4(x - 0)$$

$$\Rightarrow y = 1 + 4x.$$

Ex. Find the linear approximation for $\frac{1}{1-4(\frac{1}{e^5})}$, $\frac{1}{e^5} = 0.0067$

By above: $L(x) = 1 + 4x$

$$\text{So, } f\left(\frac{1}{e^5}\right) \approx 1 + 4\left(\frac{1}{e^5}\right) = 1 + 4(0.0067) \\ = 1.0268$$

$$\text{Actually: } f\left(\frac{1}{e^5}\right) = \frac{1}{1-4\left(\frac{1}{e^5}\right)} = 1.02769$$

Note: $\cdot \frac{d}{dx}(\log_a x) = \frac{1}{x \ln a}$, $\frac{d}{dx}[\ln x] = \frac{1}{x}$

$\cdot \log_t(ab) = \log_t a + \log_t b$

$\cdot \log_t\left(\frac{a}{b}\right) = \log_t(a) - \log_t(b)$

$\cdot \log_t(a^r) = r \log_t a$

Ex. Find $\frac{d}{dx} \left[\log_2 \left(\frac{x^2(3x+7)}{\sqrt{x^5+2}} \right) \right]$

Best Method: Use log properties first.

$$\begin{aligned} \log_2 \left(\frac{x^2(3x+7)}{\sqrt{x^5+2}} \right) &= \log_2(x^2(3x+7)) - \log_2(\sqrt{x^5+2}) \\ &= \log_2 x^2 + \log_2(3x+7) - \log_2((x^5+2)^{\frac{1}{2}}) \\ &= 2 \log_2 x + \log_2(3x+7) - \frac{1}{2} \log_2(x^5+2) \end{aligned}$$

Now, $\frac{d}{dx} \left[\log_2 \left(\frac{x^2(3x+7)}{\sqrt{x^5+2}} \right) \right]$

$$= \frac{d}{dx} \left[2 \log_2 x + \log_2(3x+7) - \frac{1}{2} \log_2(x^5+2) \right]$$

$$= 2 \frac{d}{dx} [\log_2 x] + \frac{d}{dx} [\log_2(3x+7)] - \frac{1}{2} \frac{d}{dx} [\log_2(x^5+2)]$$

$$= 2 \cdot \frac{1}{x \ln 2} + \frac{1}{(3x+7) \ln 2} \cdot 3 - \frac{1}{2} \frac{1}{(x^5+2) \ln 2} \cdot 5x^4$$

$$= \frac{2}{x \ln 2} + \frac{3}{(3x+7) \ln 2} - \frac{5x^4}{2(x^5+2) \ln 2}$$


Ex. Suppose the position vector is given $s(t) = 4t^2 - 8t + 30$, $t \geq 0$.

Find the intervals where the particle is speeding up & slowing down.

$$s(t) = 4t^2 - 8t + 30$$

$$v(t) = s'(t) = 8t - 8 \begin{cases} \text{for } t > 1, & v(t) > 0 \\ \text{for } t < 1, & v(t) < 0 \end{cases}$$

$$a(t) = s''(t) = 8, \text{ for all } t \geq 0$$



So, for $0 < t < 1$, $v(t) < 0$ & $a(t) > 0$

\Rightarrow the particle is slowing down on $(0, 1)$

for $t > 1$, $v(t) > 0$ & $a(t) > 0$

\Rightarrow the particle is speeding up on $(1, \infty)$