

Continuity:

Handwavy understanding is, it's about being able to draw graph continuously without lifting the pen/pencil from the paper.

Defⁿ 1- A function $f(x)$ is continuous at $x=a$ if & only if

- (i) $f(a)$ is defined
 - (ii) $\lim_{x \rightarrow a} f(x)$ exists
 - (iii) $\lim_{x \rightarrow a} f(x) = f(a)$
- } satisfies.

⊛ If it fails to be continuous, we say it's **discontinuous** at $x=a$.

⊛ f is continuous on an interval if f is continuous at every point a in that interval.

Determining Continuity at a point:


$$f(x) = \begin{cases} \frac{\sin x}{x}, & x \neq 0 \\ 1, & x = 0 \end{cases}$$

Then $f(0) = 1$ & we have seen $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$.

Therefore, $f(x)$ is continuous at $x=0$

Determining Continuity over an Interval:

$$f(x) = \sqrt{x-4}, \quad x \text{ is in } [4, 7)$$



Note: $[4, 7)$ means domain has the point $x=4$ in it, but not 7.

So we need to check for two cases:

Case 1: Continuity at $x=4$

We need to check for $\lim_{x \rightarrow 4^+} f(x)$ & $f(4)$ only.

$$\text{Now, } f(4) = \sqrt{4-4} = \sqrt{0} = 0$$

$$\& \lim_{x \rightarrow 4^+} f(x) = \lim_{x \rightarrow 4^+} \sqrt{x-4} = \sqrt{4-4} = 0$$

So, $\lim_{x \rightarrow 4^+} f(x)$ exists & $\lim_{x \rightarrow 4^+} f(x) = f(4)$.

$\Rightarrow f(x)$ is continuous at $x=4$.

Case 2: Continuity at any interior point $x=t$ in $[4, 7)$

$$\text{Now, } \left. \begin{aligned} \lim_{x \rightarrow t^-} f(x) &= \lim_{x \rightarrow t^-} \sqrt{x-4} = \sqrt{t-4} \\ \lim_{x \rightarrow t^+} f(x) &= \lim_{x \rightarrow t^+} \sqrt{x-4} = \sqrt{t-4} \end{aligned} \right\} \lim_{x \rightarrow t} f(x) = \sqrt{t-4}$$

& $f(t) = \sqrt{t-4}$, so $f(x)$ is cont. at any interior point of $[4, 7)$

Functions that are continuous everywhere:

① $f(x) = \text{polynomial in } x$ (say, $x^5 + 7x + 5$)

② $f(x) = |x| \rightsquigarrow$ the absolute value function.

③ $f(x) = \text{exponential functions}$ (e^x , $e^{-(x+4)}$)


④ $f(x) = \sin x / \cos x$.

⑤ $\tan x$, $\csc x$, $\sec x$, $\cot x$ are continuous on their respective domains.

$\log_a x$, $\ln x$, \sqrt{x} , $\sqrt[n]{x}$, rational functions are also continuous on their respective domains.


Discontinuities

Situation 1. $\lim_{x \rightarrow a} f(x) \text{ DNE.}$




Jump Discard.
at $x=a$


Situation 2. $\lim_{x \rightarrow a} f(x)$ exists, but not finite



Situation 3. $\lim_{x \rightarrow a} f(x)$ exists, but $f(a)$ doesn't exist.



Situation 4. Both $\lim_{x \rightarrow a} f(x)$ & $f(a)$ exists, but not equal.



So there are three type of Discontinuity:

① Infinite, when $\lim_{x \rightarrow a^\pm} f(x) = \pm \infty$

② Removable, define $f(a)$ intelligently.

③ Jump, when $\lim_{x \rightarrow a^-} f(x) \neq \lim_{x \rightarrow a^+} f(x)$

Eg. $f(x) = \frac{x^2 - 2x}{x^2 - 4} = \frac{x(x-2)}{(x+2)(x-2)}$

$$\text{Domain} = \{x \mid x \neq -2, 2\}$$

At $x = -2$,

$$\lim_{x \rightarrow -2} f(x) = \pm \infty \Rightarrow \text{Infinite discant. at } x = -2$$

At $x = 2$, $\lim_{x \rightarrow 2} f(x) = \frac{1}{2}$, but it is not defined, so $f(2)$ DNE.

\Rightarrow Removable discant at $x = 2$



Note:- If $f(x)$ is cont at $x = a$, then the limits exists at $x = a$.

But if the limit exists at $x = a$, then it might or might not be continuous at $x = a$.

Intermediate Value Theorem (IVT):

If f is continuous on $[a, b]$ [ie, you can draw a continuous curve joining $(a, f(a))$ & $(b, f(b))$], then if c is any number between $f(a)$ & $f(b)$.

Then there is at least one x_0 between a & b with $f(x_0) = c$.



Q. Is there any such value between 0 & $\pi/2$, for which $x = \cos x$ holds?

Solⁿ: Let us consider $f(x) = \cos x - x$

Note $f(x)$ is Continuous on $[0, \frac{\pi}{2}]$.

$$\text{Now, } f(\pi/2) = \cos(\pi/2) - \pi/2 = 0 - \pi/2 = -\pi/2$$

$$\text{ \& } f(0) = \cos(0) - 0 = 1 - 0 = 1$$

$$\text{Since, } -\pi/2 < 0 < 1 \Rightarrow f(\pi/2) < 0 < f(0)$$

So by IVT, there exists $\theta \in (0, \pi/2)$ such that $f(\theta) = 0$

$$\text{ie, } \cos \theta - \theta = 0 \Rightarrow \cos \theta = \theta.$$