


Linear Approximations & Differentials:



⊛ Note: Tangent is the best straight line approximation to $y = f(x)$ at $x = a$.

Now the equation of the tangent line passing through $(a, f(a))$ with slope $f'(a)$ is given by

$$y - f(a) = f'(a)(x - a)$$

$$\Rightarrow y = \underbrace{f(a) + f'(a)(x - a)}_{L(x)},$$

the linear approximation.

Ex. (a) Find the linear approximation of $f(x) = \sqrt[3]{x}$ at $a = 8$

(b) Use linear approximation to estimate $\sqrt[3]{8.05}$ & $\sqrt[3]{25}$

Solⁿ: (a) As like finding equation of tangent, we need to find

① $f(8)$ in this case &

② $f'(8)$ too.

$$\text{Now, } f(8) = \sqrt[3]{8} = 2$$

$$\begin{aligned} \& f'(x) &= [\sqrt[3]{x}]' \\ &= (x^{1/3})' = \frac{1}{3} x^{-2/3} \end{aligned}$$

$$\begin{aligned} \text{So, } f'(8) &= \frac{1}{3} (8)^{-2/3} \\ &= \frac{1}{3} (2^3)^{-2/3} \\ &= \frac{1}{3} (2^{-2}) = \frac{1}{3} \left(\frac{1}{4}\right) = \frac{1}{12} \end{aligned}$$

Therefore, the linear approximation is:


$$\begin{aligned} L(x) &= f(8) + f'(8) \cdot (x-8) \\ &= 2 + \frac{1}{12} (x-8). \end{aligned}$$

(b) To estimate $\sqrt[3]{8.05}$ & $\sqrt[3]{25}$, we replace $x = 8.05$ & $x = 25$ respectively & find out $L(8.05)$ & $L(25)$

$$\begin{aligned} \text{Now, } L(8.05) &= 2 + \frac{1}{12} (8.05 - 8) = 2 + \frac{1}{12} (0.05) \\ &= 2.00416667 \end{aligned}$$

$$\& L(25) = 2 + \frac{1}{12} (25 - 8) = 2 + \frac{17}{12} = 3.41666667$$

Note: $\sqrt[3]{8.05} = 2.00415862$ & $\sqrt[3]{25} = 2.92401774$




⊛ Remark:- For $\sqrt[3]{25}$, if we find the linear approximation around $x=25$ (say $x=27$) then we will get more accurate result.

for instance: $L(x)$ about $x=27$ is $3 + \frac{1}{27}(x-27)$

$$= 3 + \frac{x}{27} - 1$$

$$= 2 + \frac{x}{27}$$

$$\therefore L(25) = 2 + \frac{25}{27} = 2.9259259$$



Arbitrary Distance
from a

$$\Delta x = x - a$$

$$\Delta y = f(x) - f(a)$$

$$\boxed{\Delta y = f(x + \Delta x) - f(x)}$$

Infinitesimal Distance
from a .

$$dx = x - a$$

$$dy = L(x) - L(a)$$

$$= [f(a) + f'(a)(x-a)] - f(a)$$

$$= f'(a)(x-a)$$

$$= f'(a) dx$$

$$\boxed{dy = f'(x) dx} \longrightarrow \text{Differentials.}$$

Ex. Let $y = x^2 + 7$

(a) Find dy in terms of dx .

(b) Approximate Δy , if $x = 1$ & $\Delta x = 0.001$

$$dy = f'(x) dx \quad \& \quad \Delta y = f(x + \Delta x) - f(x).$$

$$(a) \quad f(x) = x^2 + 7 \Rightarrow f'(x) = 2x$$

$$\text{So, } dy = 2x dx.$$

$$(b) \quad \Delta y = f(1 + 0.001) - f(1)$$

$$= f(1.001) - f(1)$$

$$= [(1.001)^2 + 7] - [1^2 + 7]$$

$$= (1.001)^2 - 1^2 = (1.001 + 1)(1.001 - 1)$$

$$= (2.001)(0.001)$$

$$= 0.002001$$